

Occurrence and Distribution of Entomopathogenic Fungi in Cultivated Soil and Its Efficacy Against *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) Under Laboratory Conditions

Zainab Saeed^{1,2}, Usman Khalique², Farrah Zaidi¹, Muhammad Zahid², Bibi Hajra^{1,2}, Khush Bakht Latif¹, Syeda Hira Fatima^{3,4} and Syed Jawad Ahmad Shah^{2*}

¹Institute of Zoological Sciences, University of Peshawar, Pakistan

²Plant Protection Division, Nuclear Institute for Food and Agriculture (NIFA), Peshawar, Pakistan

³College of Sciences and Engineering, Flinders University, South Australia, Australia

⁴HEAL Global Research Centre, University of Canberra, ACT, Australia

Corresponding author: Syed Jawad Ahmad Shah, Plant Protection Division, Nuclear Institute for Food and Agriculture (NIFA), Peshawar, Pakistan.

Citation: Saeed, Z., Khalique, U., Zaidi, F., Zahid, M., Hajra, B., Shah, S, J, A., et al. (2026). Occurrence and Distribution of Entomopathogenic Fungi in Cultivated Soil and Its Efficacy Against *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) Under Laboratory Conditions. *Int J Biol Life Sci.* 1(1), 01-22.

Abstract

Annually, global crop harvest significantly declines due to various pest attacks. Their effective management is crucial for sustainable agricultural productivity. Native entomopathogenic fungi (EPF) have been recognized as the most promising microbiological control agents against these pests. The present study aimed to isolate locally occurring entomopathogenic fungi and assess their efficacy against the serious noctuid pest, *H. armigera*, under laboratory conditions. EPF was explored in cultivated soil from crops of two distinct agro-ecological zones (plains and foothills) in Khyber Pakhtunkhwa province, Pakistan. Using the Galleria baiting technique, fungal isolates were recovered from collected soil samples. Upon identification, these isolates belonged to 4 different EPF species, viz., *Nomuraea rileyi*, *Aspergillus parasiticus*, *A. niger*, and *A. flavus*. Results revealed that soils from the foothills exhibited a comparatively higher percentage distribution of isolates than those from the plains. *Aspergillus niger* was the most abundant fungal species in various localities and crops. The pathogenicity of four isolated species was assessed against *H. armigera* at three concentrations (1×10^6 , 1×10^7 and 1×10^8 conidia/mL). Results revealed that the *H. armigera* larvae were found to be susceptible to all tested EPF species, particularly at high concentration levels. *M. rileyi* was the most effective, causing the highest percent mortality and exhibiting the lowest percentage of pupal recovery and adult emergence, followed by *Aspergillus* species. Probit analysis showed that *M. rileyi* was highly virulent, with the lowest LC₅₀ and LT₅₀ values. This study reveals the potential of *M. rileyi* to serve as an effective biocontrol agent in

integrated pest management strategies against *H. armigera*, and as a promising candidate for bio-pesticide product development. The use of EPF agents will ensure the production of healthier organic crops by eliminating insecticide residue and resistance problems.

Keywords: Helicoverpa Armigera, Microbial Control, Entomopathogenic Fungi, Distribution, Galleria Mellonella

Introduction

The tomato fruit worm, *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae), is a highly destructive and cosmopolitan pest. Its larval stage is highly voracious and infests over 181 host plant species belonging to more than 47 families. This pest successfully adapts to various agro-ecosystems due to its polyphagous nature, facultative diapause, high reproductive capacity, and migratory potential. In Pakistan, from the late 20th century to the early 21st century, this pest was identified as a significant factor in yield reduction and threatening major field crops.

Tomatoes, *Lycopersicon esculentum* (Mill.) are among the valuable crops targeted by *H. armigera*. In Pakistan, tomatoes are the second most produced crop in the horticultural sector. The initial instars of *H. armigera* feed on green shoots and foliage, while subsequent stages target buds, inflorescences, and fruiting bodies, resulting in serious damage to post-harvest produce. This noctuid pest is estimated to cause around 53% fruit loss in tomatoes. Farmers in Pakistan and around the world mainly rely on chemical applications to control this pest. However, the excessive use of broad-spectrum insecticides has raised serious concerns about their harmful effects on human health, the environment, non-target species, disturbs ecological balance and leads to pest resurgence and the development of pest resistance. According to the Arthropod Pesticide Resistance Database (APRD) (<http://www.pesticideresistance.org/DB/>), this pest has become resistant to over 40 active ingredients. This situation has increased interest in finding more eco-friendly alternatives to reduce the harmful impacts of insect pests and support sustainable farming.

One such approach is biological control with insect-invading pathogens. Entomopathogenic fungi (EPF) are eco-friendly microbial agents naturally found in various terrestrial habitats; widely recognized worldwide in integrated pest management due to their broader host range, ability to cause lethal mycosis in insect pests through cuticle penetration, and capability to control different developmental stages more effectively than other strategies. Moreover, EPF strains promote long-term environmental sustainability by germinating on insect cadavers, thereby enhancing their inoculum density and disseminating their network throughout the ecosystem.

Metarhizium spp. are frequently used or tested as eco-friendly microbiological agents because they are easy to produce on a large scale. *Metarhizium rileyi* (Farl.), formerly known as *Nomuraea rileyi*, is a potential EPF that can cause lethal epizootics in various insects, including noctuid pests. Various studies have reported

the potential of *M. rileyi* for developing a microbial pesticide against lepidopterous pests. *M. rileyi* is widely used in control strategies for *H. armigera* due to its primary features, including environmental friendliness and host specificity. *Aspergillus* is another important genus within Ascomycota, encompassing more than 250 species. Some of these species have been identified as entomopathogenic, such as *A. niger*, *A. parasiticus*, and *A. flavus*. *Aspergillus* spp. employ a unique infection method by initiating lipid peroxidation, followed by a series of events that ultimately compromise the host's immunity. They release toxic compounds that cause gut tissue leakage and induce septicemia, leading to the death of the host insect.

Mensah et al. reported the effective control of *H. armigera* through the use of local *Aspergillus* sp. (BC 639), which also had a minimal impact on beneficial organisms. Similarly, Sebayang et al. explored native EPF species as microbial agents against *H. armigera* and isolated *A. flavus* and other EPF species from infected *H. armigera* larvae. These species have been utilized as microbial agents against other agricultural pests, including *Oligonychus coffeae*, *Dysdercus koenigii*, *Spodoptera frugiperda*, and *Spodoptera litura*. However, in Pakistan, the potential of these entomopathogens viz., *M. rileyi*, *A. niger*, *A. parasiticus*, and *A. flavus* has never been investigated against the noctuid pest, *H. armigera*.

Moreover, understanding the composition and distribution of indigenous species is vital for preserving fungal strains in soil, which is necessary for effectively managing local pest populations in agro-ecosystems. Additionally, laboratory evaluations are crucial to determine effective concentration levels, ensuring the success and safety of future field applications against this pest. The study's objectives were to explore indigenous EPF strains in cultivated soils of KP province and evaluate their potential against the local pest, *H. armigera*, using the topical inoculation method under laboratory conditions. The main parameters examined included the comparative distribution of fungal isolates in plain areas versus foothills and between agronomic versus horticultural crops, along with first mortality, total mortality (%), pupal recovery (%), adult emergence (%), and the determination of LC_{50} and LT_{50} .

Materials and Methods

Study Area

Isolation of entomopathogenic fungi and pathogenicity bioassays were performed in the Biocontrol and Plant Pathology laboratories of Nuclear Institute for Food Agriculture (NIFA), Peshawar. The morphological identification of isolated EPF species was done in the Vector Biology Laboratory of the Institute of Zoological Sciences, University of Peshawar.

Collection of Soil Samples

Soil samples were collected from four agronomic (rice, maize, sugarcane, mustard) and four horticultural crops (tomato, potato, guava, peach) from distinct geographical terrains in Khyber Pakhtunkhwa province (Plains: Peshawar and Charsadda and Foothills: Abbottabad and Haripur) during 2023-24. The details regarding the geographic coordinates of surveyed regions are given in Figure 1.




Figure 1 : Map of different agro-ecological zones of Khyber Pakhtunkhwa (KP) province, Pakistan showing soil sampling sites for entomopathogenic fungi isolation in 4 districts. The values on the X-axis present longitudinal coordinates and values on the Y-axis present latitudinal coordinates.

From each locality, five soil samples (~200 g each) were collected from a depth of 5-10 cm at five random points by using a hand shovel with three replicates per point and stored in sterile zip-lock bags (28×20 cm) at 4 °C in a refrigerator until used. The samples from each site were mixed to get a unit sample and sieved through a 16 mm mesh to remove roots, stones, litter, etc. Soil samples were then spread on rectangular trays and kept open to remove excessive moisture to avoid nematode infestation as suggested by Quesada-Moraga et al.

Isolation of Entomopathogenic Fungi

Galleria bait method was used for isolating the entomopathogenic fungi from collected soil samples. The wax moth larvae, *Galleria mellonella* (Lepidoptera: Pyralidae), were originally collected from infested bee combs acquired from local beekeepers. The *G. mellonella* culture was sustained on a suitable diet, specifically bee wax, and kept in rectangular trays within insect rearing cages (60 x 60 x 60 cm) under controlled conditions i.e., $26 \pm 2^\circ\text{C}$ temp., $65 \pm 5\%$ R.H. and continuous darkness. The dry soil samples were first moistened to field capacity, and then, after thorough mixing, from each unit sample, five sub-samples were taken into autoclaved glass vials. At the top of the soil, about 1 cm space was left and vial lids were provided with needle holes for proper aeration. Five healthy 3rd instar larvae were added to each vial using sterile forceps and incubated at $26 \pm 2^\circ\text{C}$ for ten days. The vials were inverted daily to ensure regular larval movement through the soil. After incubation, the soil was examined, and the bait larvae were removed. The

larval cadavers were first surface sterilized in 3% sodium hypochlorite (disinfectant) for 3 min, then washed with sterile distilled water for 3 min as suggested by Sookar et al. and individually incubated at $26 \pm 2^{\circ}\text{C}$ on wet filter paper in sterile glass petri plates for three days to enhance pathogen sporulation.

The cadavers with symptoms of fungal infection were transferred to Sabouraud Dextrose Yeast Agar (SDAY) plates comprising 10 g dextrose, 2.5 g peptone, 2.5 g yeast extract and 20 g agar in 1 litre of distilled water (supplemented with antibiotics 100 mg/l to inhibit bacterial growth) inside a laminar flow cabinet in Plant Pathology Laboratory, NIFA. They were kept at $26 \pm 2^{\circ}\text{C}$ temperature, $60 \pm 5\%$ relative humidity in complete darkness till the growth of fungal mycelium. The culture obtained from an infected bait larvae was considered an isolate. The name code for isolates were based according to crop: Maize (M), Sugarcane (S), Mustard (Mu), Rice (R), Potato (P), Tomato (T), Guava (G) and Peach (P); replication (R1-R3) and sample number within crop (S1-S5) and thus samples were labeled as MR1S1, RR2S5, etc.

Identification of EPF Isolates

The fungal isolates were identified based on morphological traits, such as colony appearance, spore size and shape, etc., using a taxonomic key in Vector Biology Laboratory of the Institute of Zoological Sciences, University of Peshawar. Pure culture plates of each fungal isolate were then submitted to the First Fungal Culture Bank of Lahore, Pakistan (FCBP) for confirmation. The identified fungal cultures were deposited in the Biocontrol laboratory of Nuclear Institute for Food Agriculture (NIFA), Peshawar, and Vector Biology Laboratory of the Institute of Zoological Sciences, University of Peshawar.

Insect Rearing

H. armigera larvae were collected from the NIFA field area and initially reared on a natural diet (peas) in plastic cups. Pupae were collected and transferred to glass vials for adult emergence. The emerged adults were coupled, kept in 2.5 L plastic jars covered with muslin cloth as a substrate for oviposition, and provided with a 10% sugar solution. The muslin cloth with eggs was changed daily and shifted to plastic boxes, with an artificial diet for hatchlings, comprising of chickpea powder (79.6%), scorbic acid (1%), methyl para hydroxyl benzoate (0.8%), ascorbic acid (0.3%), streptomycin (0.3%), yeast (12.7%), agar (0.1%) and vitamin B12 (0.0001%) by weight in distilled water (1000 ml). After hatching, the third instars of the F1 generation were used in bioassays. All developmental stages of *H. armigera* were kept under controlled conditions of $26 \pm 2^{\circ}\text{C}$ temperature, $65 \pm 5\%$ relative humidity, and L14: D10 hr photoperiod.

Pathogenicity of Entomopathogenic Fungi on *H. armigera*

Bioassays were performed in the Biocontrol laboratory of NIFA to evaluate the pathogenicity of isolated EPF species against *H. armigera*. Conidial suspensions were prepared from 2-3 weeks old pure EPF cultures in 10 ml of distilled water with Tween-80 (0.1%) in sterile 15 ml tubes and quantified using a Hemocytometer. The desired concentrations of each fungal strain, i.e., 1×10^8 , 1×10^7 , and 1×10^6

conidia/ml, were adjusted through serial dilution. Three replicates were used per treatment and each replicate contained nine 3rd instar of *H. armigera*, which were topically inoculated using a micropipette with a volume of 10 microliter of conidial suspensions. In a control treatment, the larvae were treated similarly, using only distilled water. The bioassay was repeated twice and during the experimental period, the larvae were provided with artificial diet and adults were fed with 10% sugar solution.

Parameters to Be Studied

Percentage Diversity of Soil-Dwelling EPF

To calculate the percentage diversity of a particular identified EPF strain (x), the following formula was used:

Total count of Identified EPF% Diversity of 'x' = No.of Identified EPF species (x) × 100

Concentration-Response Bioassay

Concentration-response tests were conducted for different parameters. In each treatment, as noted above, nine healthy third instars were placed individually in petri plates and topically inoculated with conidial suspension and in the control group, treated with distilled water. The first larval mortality data for each treatment's replication were recorded for 5 days following the inoculation of EPFs. The larvae showing no indication of movement were considered dead. The total mortality and median lethal concentration (LC₅₀) for each fungal treatment were determined at different post-inoculation intervals: total mortality after 7 and 14 days, LC₅₀ after 3, 7, and 14 days. The effect of different concentrations of EPFs on the pupal period and adult emergence of *H. armigera* was also assessed. The median lethal time (LT₅₀) was also estimated for each fungal concentration.

Research Design

A Completely Randomized Design (CRD) was followed, with 3x replicates. The aspect used was the conidial concentration treatments (T) with the following levels for each fungal species: 1×10⁶, 1×10⁷, and 1×10⁸ conidia/ml. The bioassay was conducted twice, comprising a total of thirteen treatments, including a control group.

Statistical Analysis

The bioassay data (percent mortality, pupal recovery and adult emergence) were analyzed using one-way ANOVA with Statistix version 8.1 software to determine means and different parameter values for tested fungal treatments. The means of significant treatments were compared with Tukey's HSD (Honestly Significant Difference) test at a significance level of $\alpha = 0.05$. To estimate LC₅₀ and LT₅₀ and their associated P values, degrees of freedom, chi-square and slopes of tested EPF species, probit analysis was performed on bioassay data using SPSS v16.

Results

Isolation and Identification of EPF Isolates

A total of 209 fungal isolates were recovered from baited wax moth larvae. Based on morphological traits identified using a taxonomic key and confirmed by the Fungal Culture Bank of Pakistan (FCBP), fungal isolates were classified into four different

EPF species: *Nomuraea rileyi*, *Aspergillus parasiticus*, *Aspergillus flavus*, and *Aspergillus niger*. Their morphological details are presented in Figure 2.




Figure 2: Colony visuals and microscopic view of identified EPF species (a) *Nomuraea rileyi* (b) *Aspergillus parasiticus* (c) *Aspergillus flavus* (d) *Aspergillus niger*.

Comparative Distribution of Fungal Isolates in Different Agro-Ecological Zones of KP Province

The comparative distribution of EPF isolates was determined in cultivated soil from diverse agro-ecological zones, including plains and foothills of KP Province. Of the total 209 isolates, 133 were obtained from cultivated soils in the plain area districts, i.e., Peshawar and Charsadda, while the remaining 76 isolates came from the foothills districts, i.e., Haripur and Abbottabad. Our results revealed that the comparative distribution of all isolated EPF species was higher in the soil of the foothills than in the plain areas (Figure 3). Overall, fungal species composition was uniform throughout, with the highest percentage distribution of *Aspergillus niger* i.e., 48% in the foothills and 37% in the plain regions, while *Nomuraea rileyi* had the lowest i.e., 15% in the foothills and 11% in the plain areas.




Figure 3 : illustrates the percent distribution of entomopathogenic fungi (EPF) isolates in plains vs foothills soil of Khyber Pakhtunkhwa province. The findings suggest that the comparative distribution of all EPF isolates was higher in the soil of the foothills than in the plain areas. Overall, fungal species composition was uniform throughout, with the highest percentage distribution of *Aspergillus niger*, followed by *Aspergillus flavus*, *Aspergillus parasiticus* and *Nomuraea rileyi*.

Comparative Distribution of EPF Isolates in Different Cropping Systems

The percent distribution of fungal isolates was compared across different cropping systems viz., agronomic and horticultural crops of Plains and Foothills in KP Province. Of 209 isolates, the highest number (119) was found in various horticultural crops. The percent distribution of nearly all fungal isolates in horticultural crops was relatively higher than in different agronomic crops (Table 1). Overall, the highest percent distribution was recorded for *A. niger* i.e., 43% in horticultural crops and 39% in agronomic crops, while *A. parasiticus* had the lowest percent distribution with 6% in horticultural and 5% in agronomic crops. (Table 1).

Table 1 : Effect of various cropping categories (agronomic vs horticultural) on % distribution of EPF species. The values presented reflects the number of isolates of each EPF species and its % distribution frequency.

Fungal Species	Isolates in Agronomic Crops	% distribution	Isolates in Horticultural Crops	% distribution
<i>Nomuraea rileyi</i>	18	20	10	8
<i>Aspergillus parasiticus</i>	5	6	25	21
<i>Aspergillus niger</i>	39	43	43	36
<i>Aspergillus flavus</i>	28	31	41	34

Concentration-Response Bioassays

First larval mortality of *H. armigera* exposed to different EPF concentrations

First mortality is the duration in which different EPF concentrations caused the first larval death to be recorded over 5 days. Results showed that at relatively high concentration levels (1×10^8 , 1×10^7 conidia/ml), the first larval mortality in all fungal treatments occurred in a minimum time i.e., within 2 days. *Nomuraea rileyi* was found to be the most effective, causing the first mortality in the minimum time (Table 2).

Table 2 : First mortality of *H. armigera* after being exposed to different concentrations of the tested EPF species. The sign (+) represents the first death of *H. armigera* and the sign (-) represents no death over 5 days of fungal exposure.

Treatments ^a	First Mortality of <i>H. armigera</i> (in days)				
	1 day	2 days	3 days	4 days	5 days
<i>Nomuraea rileyi</i>					
1×10^6	-	-	-	+	-
1×10^7	-	+	-	-	-
1×10^8	+	-	-	-	-
<i>Aspergillus parasiticus</i>					
1×10^6	-	-	-	-	+
1×10^7	-	+	-	-	-
1×10^8	-	+	-	-	-
<i>Aspergillus niger</i>					
1×10^6	-	-	-	-	+
1×10^7	-	+	-	-	-
1×10^8	-	+	-	-	-
<i>Aspergillus flavus</i>					
1×10^6	-	-	-	-	+
1×10^7	-	-	-	+	-
1×10^8	-	+	-	-	-
Control (water)	-	-	-	-	-

Total Percent Mortality of *H. armigera* exposed to different EPF concentrations at three post-inoculation intervals

The total mortality for each treatment was recorded at two post-inoculation intervals

(PII): 7, and 14 days against the third instar of *H. armigera*. The recorded data were analyzed using ANOVA to assess the differences among treatments and the group means were compared with Tukey's HSD (Honestly Significant Difference) test at a significance level of $\alpha = 0.05$. Our findings revealed that at 7d PII, *N. rileyi* exhibited the highest mortality percentage in the third instar at 1×10^8 conidia/ml followed by *A. niger* at the same conidial concentration, while the lowest mortality was recorded in control group ($F=13.1$, $P=0.000$). At 14d PII, the highest mortality against the third instar was recorded with higher concentrations of tested EPF and no mortality occurred in the control treatment ($F=10.1$, $P=0.000$) (Table 3). Among the tested EPF species, *N. rileyi* demonstrated the highest total mortality with 1×10^8 conidia/ml at different post-inoculation intervals.

Table 3 : Total Mortality (%) of third instar of *H. armigera* after being treated with different concentrations of the tested EPF species at two post-inoculation intervals. The values presented reflect means ($\pm$ SE), and different letters indicate statistically significant differences ($P \leq 0.05$, Tukey HSD test).

Treatments	Mortality after 7 days	Mortality after 14 days
Nomuraea rileyi		
1×10^6	44.4 BC	66.7 ABC
1×10^7	55.6 BC	100.0 A
1×10^8	100.0 A	100.0 A
Aspergillus parasiticus		
1×10^6	44.4 BC	55.6 BC
1×10^7	66.7 ABC	88.9 AB
1×10^8	55.6 BC	100.0 A
Aspergillus niger		
1×10^6	33.3 CD	44.4 C
1×10^7	44.4 BC	66.7 ABC
1×10^8	77.8 AB	100.0 A
Aspergillus flavus		
1×10^6	66.7 ABC	66.7 ABC
1×10^7	66.7 ABC	77.8 ABC
1×10^8	66.7 ABC	100.0 A
Control		

Water	0.0 D	0.0 D
Sig. Level	P=0.000	P=0.000

Effect of different EPF concentrations on Pupal Recovery (%) and Adult Emergence (%) of *H. armigera*

The statistical analysis revealed that different conidial concentrations of *N. rileyi* significantly reduced the percent pupal recovery and adult emergence of previously treated *H. armigera* larvae (Table 4). The highest percent pupal recovery was recorded in the control group, which significantly decreased with increasing concentration levels and reached its lowest at higher concentrations of 1×10^8 , 1×10^7 conidia/ml of *N. rileyi*, and 1×10^8 conidia/ml of other tested species ($F=13.1$, $P=0.000$). Similarly, the lowest percent emergence was observed in higher concentrations of the tested EPF species. While the highest percent adult emergence was recorded in the control group ($F=3.01$, $P=0.00$) (Table 4).

Table 4 : Effect of different concentrations of tested EPF species on % pupal recovery and % adult emergence of *H. armigera*. The values presented reflect means ($\pm$ SE), and different letters indicate statistically significant differences ($P \leq 0.05$, Tukey HSD test).

Treatments	Pupal recovery (%)	Adult emergence (%)
<i>Nomuraea rileyi</i>		
1×10^6	33.3 BCD	50.0 A
1×10^7	0.0 D	0.0 A
1×10^8	0.0 D	0.0 A
<i>Aspergillus parasiticus</i>		
1×10^6	44.4 BC	50.0 A
1×10^7	11.1 CD	33.3 A
1×10^8	0.0 D	0.0 A
<i>Aspergillus niger</i>		
1×10^6	55.6 B	50.0 A
1×10^7	33.3 BCD	33.3 A
1×10^8	0.0 D	0.0 A
<i>Aspergillus flavus</i>		
1×10^6	33.3 BCD	100.0 A

1×10 ⁷	22.2 BCD	66.7 A
1×10 ⁸	0.0 D	0.0 A
Control		
Water	100.0 A	100.0 A

LC₅₀ and LT₅₀ Values of the Tested EPF Species Against *H. armigera*

Using SPSS v16 and a probit analysis on mortality data, the LC₅₀ and LT₅₀ values were calculated for the EPF species, setting P<0.05 as the significance level. We measured LC₅₀ values at three post-inoculation intervals: 3, 7, and 14 days, against the third instar of *H. armigera*. Our findings showed that the LC₅₀ values of the tested EPF species were time-dependent and decreased significantly as the post-inoculation interval increased. At 3d, the lowest LC₅₀ was recorded for *N. rileyi* (1.36×10⁸ spores/ml), followed by *A. niger*, *A. flavus* (1.15×10⁹ spores/ml) and *Aspergillus parasiticus* (1.66×10⁹ spores/ml) (Table 5). At 7d, *N. rileyi* was the most virulent, with the lowest LC₅₀ (2.40×10⁶ spores/ml), followed by *A. parasiticus* (3.14×10⁶ spores/ml) and *A. niger* (8.06×10⁶ spores/ml), while *A. flavus* was the least effective, showing the highest LC₅₀ (3.18×10⁷ spores/ml) (Table 6). At 14d, the lowest LC₅₀ was recorded for *A. flavus* (4.00×10⁵ spores/ml) followed by *N. rileyi* (7.19×10⁵ spores/ml), *A. parasiticus* (8.05×10⁵ spores/ml) and *A. niger* (1.88×10⁶ spores/ml) (Table 7).

Table 5. LC₅₀ values (conidia/ml) of entomopathogenic fungi against third instar of *H. armigera* after 3 days.

Fungal Species	N	LC ₅₀ * (conidia/mL)	Slope ± SE	Intercept	X ² (df=1)	P
<i>Nomuraea rileyi</i>	27	1.36×10 ⁸	0.52 ± 0.34	-4.20	0.20	0.65b
<i>Aspergillus parasiticus</i>	27	1.66×10 ⁹	0.43 ± 0.36	-3.94	0.36	0.55b
<i>Aspergillus niger</i>	27	1.15×10 ⁹	0.39 ± 0.34	-3.53	0.01	0.92b
<i>Aspergillus flavus</i>	27	1.15×10 ⁹	0.39 ± 0.34	-3.53	0.01	0.92b

N=number of treated insects, SE=Standard error, X² =Chi square value, df=degree of freedom, P=Probability value.

Table 6. LC₅₀ values (conidia/ml) of entomopathogenic fungi against third instar of *H. armigera* after 7 days.

Fungal Species	N	LC ₅₀ * (conidia/mL)	Slope SE	± Intercept	χ ² (df=1)	P
Nomuraea rileyi	27	2.40×10 ⁶	0.94 0.39	±-5.96	2.08	0.15b
Aspergillus parasiticus	27	3.14×10 ⁶	0.28 0.30	±-1.85	0.00	0.99b
Aspergillus niger	27	8.06×10 ⁶	0.59 0.32	±-4.08	0.34	0.56b
Aspergillus flavus	27	3.18×10 ⁷	0.28 0.30	±-2.14	0.00	0.99b

N=number of treated insects, SE=Standard error, X² =Chi square value, df=degree of freedom, P=Probability value.

Table 7. LC₅₀ values (conidia/ml) of entomopathogenic fungi against third instar of *H. armigera* after 14 days.

Fungal Species	N	LC ₅₀ * (conidia/mL)	Slope SE	± Intercept	χ ² (df=1)	P
Nomuraea rileyi	27	7.19×10 ⁵	2.97 5.10	±-17.38	0.00	0.95b
Aspergillus parasiticus	27	8.05×10 ⁵	1.20 0.58	±-7.10	0.08	0.77b
Aspergillus niger	27	1.88×10 ⁶	1.02 0.42	±-6.40	0.02	0.31b
Aspergillus flavus	27	4.00×10 ⁵	0.77 0.44	±-4.31	0.86	0.35b

N=number of treated insects, SE=Standard error, X² =Chi square value, df=degree of freedom, P=Probability value.

Median Lethal Time (LT₅₀)

Median lethal time (LT₅₀) was also estimated for each fungal concentration, i.e., 1×10⁶, 1×10⁷, and 1×10⁸ conidia/ml. At 1×10⁶ conidia/ml, the lowest LT₅₀ was observed in *A. parasiticus* (8.77 days), followed by *A. flavus* (9.62 days), *M. rileyi* (9.72 days), and *A. niger* (12.9 days) (Figure 4 a-c). At 1×10⁷ conidia/ml, *M. rileyi* showed the lowest LT₅₀, followed by *A. flavus* and *A. parasiticus*, with *A. niger* having the highest LT₅₀ (10.6 days) (Figure 4 b). Similarly, at 1×10⁸ conidia/ml, *M. rileyi* was the most virulent with the lowest LT₅₀ (5.45 days), followed by *A. niger* (8.25 days) and *A. parasiticus* (8.49 days), while the highest LT₅₀ was recorded for *A. flavus* (8.55 days) (Figure 4 c).






Figure 4 : a-c: LT₅₀ (days) at three conidial concentrations (a) 1×10^6 conidia/ml (b) 1×10^7 conidia/ml (c) 1×10^8 conidia/ml of entomopathogenic fungi (EPF) to kill 50% population of the exposed *H. armigera* larvae. The results suggest that with increasing conidial concentration of EPF species (except *A. parasiticus*), the LT₅₀ values significantly decrease, with the minimum values recorded at 1×10^8 conidia/mL of tested EPF species.

Discussion

Many studies have reported the isolation and identification of fungal entomopathogens in different habitats. However, in Pakistan, due to limited knowledge about the occurrence and distribution of native fungal species, no microbial pesticide has been registered or made commercially available to farmers for local pest control. This study confirms the presence of potential EPF in cultivated soils. Our results revealed that the distribution percentage of fungal isolates was higher in the soil of the foothills compared to the plains, indicating a significant presence in the soils of the higher altitude districts (Haripur and Abbottabad). These findings align with those of Wakil et al., who evaluated the occurrence of fungal entomopathogens in different regions of Punjab province and reported the highest distribution of fungal isolates in soil from higher altitudes (above 600 m). While Liu et al. identified ultraviolet exposure as a major factor negatively affecting the occurrence and persistence of EPF species in soil, which contradicts our findings, as foothill soils, being at higher altitudes, are more exposed to ultraviolet radiation than those in plain areas. The difference may be due to variations in the organic content and soil texture that significantly influence the occurrence of EPF in soil habitats.

The comparative distribution of fungal isolates was higher in different horticultural crops than in agronomic fields. Our results are in partial agreement with Qayyum et al., who reported higher fungal distribution frequency in fruits (52.50%) and vegetables (42.50%) than in field crops (37.50%). Similarly, Sookar et al. also observed that the occurrence and distribution of EPF were significantly influenced by habitat type. They recovered a larger number of *Metarhizium anisopliae* isolates from the vegetable soil than from other plantations. Notably, in our results, *Aspergillus* was the most prevalent genus, with *Aspergillus niger* found as the most abundant EPF species among the fungal isolates, followed by *A. flavus*. Our results are supported by Qayyum et al., who reported *A. niger* as the most abundant fungal strain with a distribution frequency of 27.50% followed by *A. flavus* with 22.50% occurrence in sampled soil. This is because various ecological and environmental factors influence the distribution and abundance of EPF in the soil habitat. Similarly, Wakil et al. recovered *Aspergillus* as the most common genus from soil samples.

The virulence of isolated EPF species, identified as *N. rileyi*, *A. parasiticus*, *A. niger*, and *A. flavus* was evaluated in bioassays against the third instar of *H. armigera*. The larvae were found susceptible to all tested fungal entomopathogens, resulting in significant mortality, especially at their highest concentration 1×10^8 conidia/ml. Our results showed a dose-dependent mortality response, with *N. rileyi* as the most effective treatment, causing the highest percentage mortality, followed by *Aspergillus* species. These findings are partly consistent with those of Hazarika et

al., who reported concentration-dependent mortality with *N. rileyi* and identified it as a promising biocontrol agent, causing 86% mortality in *H. armigera* at a concentration of 1×10^9 spores/mL. The difference in the maximum mortality recorded in the present study and Hazarika et al. is attributed to variations in strains and the concentrations assessed in both studies. Our results are highly consistent with those of Dev et al., who found *M. rileyi* to be highly virulent against *H. armigera*, with the highest mortality observed in the third instar at 1×10^8 conidia/mL. *M. rileyi* causes host-preferential epizootics in susceptible larvae mainly by blocking respiratory structures, leading to host death. Similarly, Ramos et al. recorded an effective mortality rate (70 to 98.7%) in *Spodoptera frugiperda* larvae when treated with 24 different isolates of *M. rileyi* in bioassays.

Several studies have reported the entomopathogenic potential of *Aspergillus* species against various insect pests, including lepidopterans, with limited research focused on *H. armigera*. Mensah et al. assessed the efficacy of *Aspergillus* sp. (BC 639) against *Helicoverpa* spp. and predatory insects both in vitro and in cotton fields. They reported effective pest control using *Aspergillus* spp. with minimal effects on beneficial organisms, maintaining crop yield. Similarly, Kaur et al. reported a 63% larval mortality in *Spodoptera litura* using ethyl acetate extract of *A. niger* compared to 3% in the control group.

The pupal recovery and adult emergence of treated *H. armigera* larvae were significantly reduced with increasing conidial concentration. The lowest percent pupal recovery and percent emergence were observed at different concentrations of *N. rileyi*, followed by *Aspergillus* species. These findings partially agree with those of Mandoukas, who reported concentration-dependent pupation and emergence when exposing *H. armigera* larvae to different concentrations of *Metarhizium robertsii*, with the lowest pupation and emergence observed at the highest concentration (1×10^7 conidia/mL). However, Shanthakumar et al. (2010) found no significant effect of EPF concentration on pupation and recorded 90% pupal recovery in *S. litura* at both lower and higher concentrations of *N. rileyi* (10^4 and 10^7 conidia/mL). Similarly, Hatting et al. observed accelerated pupation in *H. armigera* after treating the larvae with *N. rileyi* using both topical and oral inoculation methods. The observed discrepancies in pupal recovery results may be attributed to the genetic makeup of fungal strains isolated from different geographical regions. Additionally, variations in methodology, including inoculation techniques and the susceptibility of targeted pests to fungal infections, also explain the contrasting outcomes.

Probit analysis revealed that the estimated LC_{50} values of the tested EPF species were time-dependent and significantly decreased as the PII increased. *N. rileyi* was found to be most virulent, exhibiting the lowest LC_{50} at 7d PII, followed by *A. parasiticus*, *A. niger* and *A. flavus*. These results are in accordance with those of Liu et al., who reported an LC_{50} of 6.24×10^6 conidia/mL for *M. rileyi* against third-instar *S. litura*. Similarly, Ramos et al. documented the LC_{50} range (2.04×10^5 to 1.05×10^6 conidia/mL) of 5 different *N. rileyi* isolates against *S. frugiperda* larvae. However, LC_{50} results for *N. rileyi* are in partial agreement with those of Dev et al., who

determined the LC_{50} of 2.81×10^5 spores/ml for *M. rileyi* against *H. armigera* third instar. This partial variation in the findings may be attributed to the genotypic variation, related to their adaptation to different native environments.

LT_{50} values of almost all fungal treatments (except *A. parasiticus*) were dose-dependent and significantly decreased with increasing conidial concentrations. These findings are supported by Yang et al., who also observed concentration-dependent LT_{50} values of *M. rileyi* against *S. frugiperda* larvae. Contrary to these results, Ramos et al. recorded LT_{50} values for 1×10^8 conidia/ml concentration of five *M. rileyi* isolates ranging from 7.04 days to 9.46 days against the second instar of *S. frugiperda*. The observed discrepancies in the LT_{50} results may be attributed to experimental conditions and targeted lepidopterous species.

Based on our findings, the distribution of entomopathogenic fungi in the different agro-ecological zones and cropping systems was not uniform throughout the soil habitat. EPF species viz.,

N. rileyi, *A. parasiticus*, *A. flavus* and *A. niger*, isolated from cultivated soil in Khyber Pakhtunkhwa province of Pakistan, effectively managed the local pest, *H. armigera* in laboratory conditions. Among all the fungal species tested, *N. rileyi* showed the most promising control, followed by *Aspergillus* species. *N. rileyi* caused deadly mycosis in *H. armigera* and showed a detrimental impact on its biological parameters as well. Hence, it is potentially suitable for the eco-friendly management of this pest and would be a sustainable alternative to synthetic insecticides. Naturally occurring EPF causes deadly mycosis in insect pests, with the potential to infect all their life stages. Their incorporation into IPM strategies will ensure the production of healthier organic crops by addressing the insecticide residue and resistance issues. Further, field trials using the extracts of isolated EPF species against *H. armigera* are needed [1-76].

Author Contributions: Zainab Saeed (Conceptualization [equal], Writing—original draft [equal], Formal—analysis and data curation [equal], Methodology [equal], Project Management, Writing—review and editing [equal]), Usman Khalique (Investigation, Formal analysis [equal], Writing—review and editing [equal], Resources, Software [equal]), Farrah Zaidi (Supervision, Formal analysis [equal], Writing—review and editing [equal], Resources [equal]), Muhammad Zahid (Writing—review and editing [equal]), Bibi Hajra (Writing—review and editing [equal]), Khush Bakht Latif (Writing—review and editing [equal]), Syeda Hira Fatima (Writing—review and editing, [equal], Map—Assistance), and Syed Jawad Ahmad Shah (Supervision, Writing—review and editing [equal], Resources [equal]).

Data Availability Statement: All data supporting the findings of this study are included in the manuscript. The raw data can be made available by the corresponding author upon reasonable request.

Acknowledgments: The authors would like to thank the Nuclear Institute for Food and Agriculture (NIFA), Peshawar and the Institute of Zoological Sciences, University of Peshawar for their support and facilitation of this research.

Conflicts of Interest: All authors declare no conflict of interest.

References

1. Ai, X., Wei, Y., Huang, L., et al. (2018). Developmental control of *Helicoverpa armigera* by ingestion of bacteria expressing dsRNA targeting an arginine kinase gene. *Biocontrol Sci. Technol*, 28(3), 253-267.
2. Askar, M., Chen, C., Borham, A., et al. (2024). First activity of *Aspergillus flavus* as entomopathogen infecting *Diaphania indica* and control efficacy across different developmental stages. *Horticulturae*, 10(12), 1315.
3. Bidochka, M. J., Kamp, A. M., Lavender, T. M., Dekoning, J., and De Croos, J. N. A. (2001). Habitat association in two genetic groups of the insect-pathogenic fungus *Metarhizium anisopliae*: Uncovering cryptic species. *Appl. Environ. Microbiol*, 67, 1335-1342.
4. Bing, D. S. and Xing, Z. L. (2008). Occurrence and diversity of insect-associated fungi in natural soils in China. *Appl. Soil Ecol*, 39, 100-108.
5. Bird, L. J. (2017). Genetics, cross-resistance and synergism of indoxacarb resistance in *Helicoverpa armigera* (Lepidoptera: Noctuidae). *Pest Manag. Sci*, 73, 575–581.
6. Chaitanya, T., Sreedevi, K., Krishna, T., et al. (2014). Biology and population dynamics of *Helicoverpa armigera* (Hubner) in *Cajanus cajan* (L.) Millsp. *Ann. Pl. Protec. Sci*, 22, 287-290.
7. Chen, C., Han, P., Yan, W., et al. (2018). Uptake of quercetin reduces larval sensitivity to lambda-cyhalothrin in *Helicoverpa armigera*. *J. Pest Sci*, 91(1), 919-926.
8. Chohan, T. Z. and Ahmad, S. (2008). An assessment of tomato production practices in Danna Katchely, Azad Jammu Kashmir. *Pak J Life Soc. Sci*, 6, 96–102.
9. Couceiro, J. C., De Fine Licht, H. H., Delalibera, I. Jr., and Meyling, N. V. (2022). Comparative gene expression and genomics reflect geographical divergence in the plant symbiotic and entomopathogenic fungal genus *Metarhizium*. *Fungal Ecol*, 60, 101190.
10. Czepak, C., Albernaz, K. C., Vivan LM., et al. (2013). First reported occurrence of *Helicoverpa armigera* (Hubner) (Lepidoptera: Noctuidae) in Brazil. *Pesqui. Agropecu. Trop*, 43, 110-113.
11. David, B. V. (2008). Biotechnological approaches in IPM and their impact on environment. *J. Biopestic*, 1(1), 3-5.
12. Dev, B., Verma, S. C., Sharma P.L, et al. (2021). Evaluation of *Metarhizium rileyi* Farlow (Samson) impregnated with azadirachtin and indoxacarb against *Helicoverpa armigera* (Hubner). *Egypt J Biol Pest Control*, 31, 142.
13. Dinesh, K., Anusha, S., Singh, R. B., et al. (2017). Estimation of avoidable yield losses caused by *Helicoverpa armigera* (Hubner) on chickpea. *J. Entomol. Zool. Stud*, 5, 1476-1478.
14. Fathipour, Y., Baghery, F., Bagheri, A., et al. (2020). Development, reproduction and life table parameters of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on five main host plants. *J. Crop Prot*, 9(4), 551-561.
15. Fite, T., Tefera, T., Negeri, M., et al. (2019). Farmer's status, knowledge & management practices on major chickpea insect pests in some selected zones of

- Ethiopia. *J. Agric. Sci*, 11(1), 31–46.
16. Fitriana, Y., Suharjo, R., Swibawa¹, I. G., et al. (2021). *Aspergillus oryzae* and *Beauveria bassiana* as entomopathogenic fungi of *Spodoptera litura* Fabricius (Lepidoptera: Noctuidae) infesting corn in Lampung, Indonesia. *Egypt. J. Biol. Pest. Control*, 31, 127.
 17. Gorczyca, A., Galon, A., Tatoj, A., et al. (2018). Occurrence of entomopathogenic fungi in agricultural and natural soils in south-eastern Poland. *J. Res. Appl. Agric. Eng*, 63(2), 63-67.
 18. Greenfeld, B. P., Peace, A., Evans, H., et al. (2015). Identification of *Metarhizium* strains highly effective against *Aedes*, *Anopheles*, and *Culex* larvae. *Biocontrol Sci. Technol*, 25, 487-502.
 19. Khan, H. A. A., Khan, T., and Iqbal, N. (2025). Biocontrol potential of isolates of *Aspergillus flavus* and *Aspergillus parasiticus* against *Rhyzopertha dominica*, *Trogoderma granarium*, and *Tribolium castaneum*. *J. Stored Prod. Res*, 114, 102799.
 20. Hatting, J. L. (2012). Comparison of three entomopathogenic fungi against the bollworm, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae), employing topical vs per os inoculation techniques. *Afr. Entomol*, 20(1), 91-100.
 21. Hazarika, S., Patgiri, P., Dutta, P., et al. (2016). Efficacy of local isolate of *Nomuraea rileyi* (Farlow) Sampson against *Helicoverpa armigera* (Hubner). *J. Entomol. Zool. Stud*, 4(3), 167-169.
 22. Humber, R. A. (2005). Entomopathogenic fungal identification. *USDA-ARS Biological Integrated Pest Management Laboratory*, 1-32.
 23. Hussein, K. A., Abdel-Rahman, M. A. A., Abdel-Mallek, A. Y., et al. (2010). Climatic factors interference with the occurrence of *Beauveria bassiana* and *Metarhizium anisopliae* in cultivated soil. *Afr. J. Biotechnol*, 9, 7674–7682.
 24. Idrees, A., Qadir, Z. A., Akutse, K. S., et al. (2021). Effectiveness of entomopathogenic fungi on immature stages and feeding performance of fall armyworm, *Spodoptera frugiperda* (Lepidoptera: Noctuidae) larvae. *Insects*, 12, 1044.
 25. Ingle, Y., Sable, Y., and Parate, R. (2017). Pathogenic and extracellular enzymes variability in entomopathogenic fungus *Nomuraea rileyi* (Farlow) Samson. *The Bioscan*, 12, 1475-1478.
 26. Iqbal, M., Gogi, M. D., Atta, B., et al. (2021). Assessment of pathogenicity of *Beauveria bassiana*, *Metarhizium anisopliae*, *Verticillium lecanii* and *Bacillus thuringiensis* var. *kurstaki* against *Bactrocera cucurbitae* Coquillett (Diptera: Tephritidae) via diet-bioassay technique under controlled conditions. *Int. J. Trop. Insect Sci*, 41, 1129–1145.
 27. Jaronski, S. T. (2010). Ecological factors in the inundative use of fungal entomopathogens. *BioControl*, 55(1), 159-185.
 28. Jones, C. M., Parry, H., Tay, W. T., et al. (2019). Movement ecology of pest *Helicoverpa*: implications for ongoing spread. *Annu. Rev. Entomol*, 64, 277–295.
 29. Karim, S. (2000). Management of *Helicoverpa armigera*: a review and prospectus for Pakistan. *Pak. J. Biol. Sci*, 3(8), 1213-1222.
 30. Karthi, S., Vasantha-Srinivasan, P., Senthil-Nathan, S., et al. (2024). Entomopathogenic fungi promising biocontrol agents for managing lepidopteran pests: Review of current knowledge. *Biocatal. Agric. Biotechnol*, 58, 103146.

31. Katsikis, C. I., Wang, P., and Zalucki, M. P. (2020). Life history traits of a key agricultural pest, *Helicoverpa armigera* (Lepidoptera: Noctuidae): are laboratory settings appropriate. *Austral. Entomol*, 59(1), 189–201.
32. Kaur, T., Datta, R., Mahajan, A., et al. (2025). Endophytic colonization of cauliflower by *Aspergillus flavus* and *Aspergillus niger* reduces performance and host preference in tobacco caterpillar, *Spodoptera litura*. *Entomol. Exp. Appl.*
33. Kaur, T., Kaur, J., Kaur, A., et al. (2016). Larvicidal and growth inhibitory effects of endophytic *Aspergillus niger* on a polyphagous pest, *Spodoptera litura*. *Phytoparasitica*, 44, 465-476.
34. Kepler, R. M., Humber, R. A., Bischof, J. F., et al. (2014). Clarification of generic "ans" species boundaries for *Metarhizium* and related fungi through multigene phylogenetics. *Mycologia*, 106, 824.
35. Kotta-Loizou, I. (2021). Diversity of mycoviruses in Aspergilli. In D. H. Bamford and M. Zuckerman (Eds.), *Encyclopedia of Virology* (4th ed.). *Academic Press*, 450-456.
36. Kriticos, D. J., Ota, N., Hutchison, W. D., et al. (2015). The potential distribution of invading *Helicoverpa armigera* in North America: Is it just a matter of time?. *PLoS ONE*, 10(3), e0119618.
37. Kumari, S., Verma, D., and Kumar, D. (2019). Evaluation of *Aspergillus niger* as a biocontrol Agent in the insect pest management of red cotton bug, *Dysdercus koenigii* (Heteroptera: Pyrrhocoridae). *J. Sci. Res*, 11(2), 235-247.
38. Liu, Y. C., Ni, N. T., Chang, J. C., et al. (2021). Isolation and selection of entomopathogenic fungi from soil samples and evaluation of fungal virulence against insect pests. *J. Vis. Exp*, 175, e62882.
39. Liu, S., Xu, Z., Wang, X., Zhao, L., Wang, G., Li, X., and Zhang, L. (2019). Pathogenicity and in vivo development of *Metarhizium rileyi* against *Spodoptera litura* (Lepidoptera: Noctuidae) larvae. *Journal of Economic Entomology*, 112(4), 1598-1603.
40. Mahankuda, B. and Bhatt, B. (2019). Potentialities entomopathogenic fungus *Beauveria bassiana* as a biocontrol agent: A review. *J. Entomol. Zool. Stud*, 7(5), 870-874.
41. Maina, U. M., Galadima, I. B., Gambo, F. M., et al. (2018). A review on the use of entomopathogenic fungi in the management of insect pests of field crops. *J. Entomol Zool Stud*, 6(1), 27-32.
42. Mantzoukas, S., Kitsiou, F., Lagogiannis, I., et al. (2022). Potential use of *Fusarium* isolates as a biological control agent against *H. armigera* (Hubner) (Lepidoptera: Noctuidae) case study. *Appl. Sci*, 12, 8918.
43. Mantzoukas, S. (2019). The effect of *Metarhizium robertsii* and *Bacillus thuringiensis* against *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae). *Adv. Ecol. Environ. Res*, 4, 136-146.
44. Mazid, S., Rajkhowa, R. C., and Kalita, J. C. (2015). Pathogenicity of *Aspergillus Niger* and *Aspergillus flavus* on red spider mite (*Oligonychus coffeae* Nietner), a serious pest of tea. *J. Entomol Zool Stud*, 3(3), 11-13.
45. Mensah, R. K., Young, A., and Rood-England, L. (2015). Development of a Microbial-Based Integrated Pest Management Program for *Helicoverpa* spp. (Lepidoptera: Noctuidae) and Beneficial Insects on Conventional Cotton Crops in Australia. *Insects*, 6(2), 333-351.

46. Meyling, N. V. and Eilenberg, J. (2006). Occurrence and distribution of soil borne entomopathogenic fungi within a single organic agroecosystem. *Agr. Ecosyst. Environ*, 113(1-4), 336-341.
47. Mishra, G. and Omkar. (2021). Gram pod borer (*Helicoverpa armigera*). In Polyphagous Pests of Crops. *Springer, Singapore*, 311-348.
48. Moore, D., Lord, J. C., and Smith, S. M. (2012). Pathogens. In B. Subramanyam and D. W. Hagstrum (Eds.), *Alternatives to Pesticides in Stored-Product IPM. Kluwer Academic Press, Dordrecht, The Netherlands*, 193-227.
49. Nnakumusana, E. S. (1985). Laboratory infection of mosquito larvae by entomopathogenic fungi with particular reference to *Aspergillus parasiticus* and its effects on fecundity and longevity of mosquitoes exposed to conidial infections in larval stages. *Current Science*, 54(23), 1221-1228.
50. Pratisoli, D., Pirovani, V. D., and Lima, W. L. (2015). Occurrence of *Helicoverpa armigera* (Lepidoptera: Noctuidae) on tomato in the Espírito Santo state. *Hortic. Bras*, 33, 101-105.
51. Qayyum, M. A., Saeed, S., Wakil, W., et al. (2021). Diversity and correlation of entomopathogenic and associated fungi with soil factors. *J. King Saud Univ. Sci*, 33(1), 101520.
52. Quesada-Moraga, E., Navas-Cortes, J. A., Maranhao, E. A. A., et al. (2007). Factors affecting the occurrence and distribution of entomopathogenic fungi in natural and cultivated soils. *Mycol. Res*, 111, 947-966.
53. Ramos, Y., Pineda-Guillermo, S., Tamez-Guerra, P., et al. (2024). Natural prevalence, molecular characteristics, and biological activity of *Metarhizium rileyi* (Farlow) isolated from *Spodoptera frugiperda* (J. E. Smith) larvae in Mexico. *J. Fungi*, 10, 416.
54. Riaz, S., Johnson, J. B., Ahmad, M., et al. (2021). A review on biological interactions and management of the cotton bollworm, *Helicoverpa armigera* (Lepidoptera: Noctuidae). *J. Appl. Entomol*, 145, 467-498.
55. Sabbour, M. M. and Abdel-Rahman, A. (2013). Efficacy of isolated *Nomuraea rileyi* and Spinosad against corn pests under laboratory and field conditions in Egypt. *Annu. Rev. Res. Biol*, 3(4), 903-912.
56. Saljoqi, A. U. R., Amin, S., Salim, M., et al. (2022). Pesticide residue analysis of three different pesticides used against *Helicoverpa armigera* (Hubner) in tomato crop. *Sarhad J. Agric*, 38(2), 448-455.
57. Sanchez-Peña, S. R. (2000). Entomopathogens from two Chihuahuan desert localities in Mexico. *BioControl*, 45, 63-78.
58. Sebayang, A., Tenrirawe, A., and Mirsam, H. (2021). Exploration of entomopathogenic fungi as potential biocontrol of corn earworm *Helicoverpa armigera* (Hübner). *IOP Conf. Ser.: Earth Environ. Sci*, 911, 012064.
59. Sevim, A., Demir, I., Hofte, M., et al. (2010). Isolation and characterization of entomopathogenic fungi from hazelnut-growing region of Turkey. *BioControl*, 55, 279-297.
60. Sezen, K. and Demirbag, Z. (2006). A new isolate of *Melolontha melolontha* entomopoxvirus in Turkey: morphology, infectivity and prevalence in the field. *Appl. Entomol. Zool*, 41, 471-477.
61. Shah, J. A., Inayatullah, M., Sohail, K., et al. (2008). Efficacy of botanical extracts and a chemical pesticide against tomato fruit worm, *Helicoverpa* Shaheen, N. *Is*

- organic farming a suitable solution for Pakistan? SDPI Res. News Bull*, 15, 78-81.
62. Shanthakumar, S. P., Murali, P. D., Malarvannan, S., et al. (2010). Laboratory evaluation on the potential of entomopathogenic fungi, *Nomuraea rileyi*, against tobacco caterpillar, *Spodoptera litura* Fabricius (Noctuidae: Lepidoptera) and its safety to *Trichogramma* sp. *J. Biopestic*, 3(1), 132-137.
 63. Sinha, K. K., Choudhary, A. K., and Kumari, P. (2016). Eco-friendly pest management for food security: entomopathogenic fungi. *Academic Press*, 475-505.
 64. Sookar, P., Bhagwant, S., Ouna, and EA (2008). Isolation of entomopathogenic fungi from the soil and their pathogenicity to two fruit fly species (Diptera: Tephritidae). *J. Appl. Entomol*, 132, 778-788.
 65. Sorathiya, K., Kalariya, S., and Patel, L. (2023). *Metarhizium anisopliae* as an entomopathogenic fungi: optimization of mass production with diverse grain substrates. *IJMCE*, 5(4), 87-95.
 66. Sun, Z., Xu, C., Chen, S., et al. (2019). Exposure to herbicides prime P450-mediated detoxification of *Helicoverpa armigera* against insecticide and fungal toxin. *Insects*, 10(1), 28.
 67. Tkaczuk, C., Król, A., Majchrowska-Safaryan, A., et al. (2014). The occurrence of entomopathogenic fungi in soils from fields cultivated in a conventional and organic system. *J. Ecol. Eng*, 15(4), 137-144.
 68. Uzman, D., Pliester, J., Leyer, I., et al. (2019). Drivers of entomopathogenic fungi in organic and conventional vineyard soils. *Appl. Soil. Ecol*, 133, 89-97.
 69. Vega, F. E., Goettel, M. S., Blackwell, M., et al. (2009). Fungal entomopathogens: New insights on their ecology. *Fungal Ecol*, 2(4), 149-159.
 70. Vega, F. E., Meyling, N. V., Luangsa-ard, J. J., et al. (2012). Fungal entomopathogens. *Insect Pathol*, 2, 171-220.
 71. Vega-Aquino, P. and Sanchez-Peña S. Blanco, C. A. (2010). Activity of oil-formulated conidia of the fungal entomopathogens *Nomuraea rileyi* and *Isaria tenuipes* against lepidopterous larvae. *J. Invertebr. Pathol*, 103, 145-149.
 72. Wakil, W., Ghazanfar, M. U., Riasat, T., et al. (2013). Occurrence and diversity of entomopathogenic fungi in cultivated and uncultivated soils in Pakistan. *Entomol. Res*, 43, 70-78.
 73. Wang, N. C. and Li, Z. H. (1984). Studies on the biology of cotton bollworm (*Heliothis armigera* Hubner) and tobacco budworm (*Heliothis assulta* Quenee). *J. Shandong Agr. Univ*, 1-2(1), 13-25.
 74. Yang, X., Zhang, Y., Zhou, J., et al. (2024). Pathogenicity, infection process, physiological and biochemical effects of *Metarhizium rileyi* against *Spodoptera frugiperda* (J. E. Smith) (Lepidoptera: Noctuidae) larvae. *Egypt J Biol Pest Control*, 34(19), 1-9.
 75. Yuan and Yong, S. (2010). Infection effect of *Nomuraea rileyi* MZ060806-XW against *Helicoverpa armigera* larvae. *J. Anhui. Agri. Sci*, 22, 11918-11999.
 76. Zimmermann, G. (1986). The "Galleria bait method" for detection of entomopathogenic fungi in soil. *J. Appl. Entomol*, 102, 213-215.